IUI Injectable Teaching Packet

OVULATION INDUCTION / (Intrauterine Insemination) "IUI" PROGRAM

Center for Advanced Reproductive Services

FARMINGTON OFFICE: 844-467-3483 or 860-321-7082

HARTFORD OFFICE: 860-525-8283

Welcome to the On-Line IUI Injectable Teaching Program at The Center for Advanced Reproductive Services. Before starting your injectable medications, the following steps MUST be completed:

- 1. All medical pre-requisites prescribed by your physician must be completed. This includes update blood work, physical exams, HSGs etc.
- 2. Your insurance pre-authorizations <u>must</u> be completed with your Financial Service Representative (FSR). You <u>cannot</u> start a cycle until "financially cleared".
- 3. Once above completed, call your primary ("clinical") RN to obtain your medication plan and a list of medication videos that you will need to watch on-line through our website. (See "MEDICATION CHECKLIST ADDENDUM for list and website instructions)
- 4. You are responsible for thoroughly reading this entire (Ovulation Induction) IUI Injectable Teaching Packet
- 5. You are responsible for reviewing all assigned medication videos. See "MEDICATION CHECKLIST" ADDENDUM at the end of the packet.
- 6. Once you have reviewed both this teaching packet AND the medication videos assigned by your primary ("clinical") nurse, call your clinic nurse and leave a message that you have done so and are ready to begin your treatment.
- 7. Your call will be returned WITHIN A FEW DAYS to complete a short phone interview to verify your understanding of your current medication protocol; to answer any questions that you might have; and to receive your actual prescription.
- 8. Consents MUST be signed BEFORE you may start medications. Please note: The consents must be witnessed by an employee of our Center (not an acquaintance or relative) OR must be signed by a public notary.
- 9. Once all the above is completed, you will be able to start your cycle.

TABLE OF CONTENTS

Part 1	About our Ovulation Induction Program
Part 2	Steps Involved in a Cycle
Part 3	Getting Started with your Cycle
Part 4	Time to Review the Website Videos

Addendums:

- 1. Helpful Hints for Injection Techniques
 Helpful Hints for Crinone Progesterone
 Helpful Hints for Endometrin Progesterone
 Helpful Hints for Progesterone Suppositories.
- 2. Medication Checklist
- 3. Injection Site Illustration
- 4. "Gonadotrophin" flow sheet (your medication plan)
- 5. Menopur visual instructions for use <u>without</u> Q cap. (same technique can also be used to mix "Luveris" medication)
- 6. "Superovulation" and "Intrauterine Inseminations (IUI)" Consents
- 7. On- Line Patient Education Questionnaire (please return this to us- we appreciate your thoughts and suggestions)

PART 1 ABOUT OUR OVULATION INDUCTION/IUI PROGRAM

IMPORTANT INFORMATION FOR YOU TO KNOW BEFORE AND DURING YOUR CYCLE

WORKING WITH THE TEAM

The doctors, nurses, laboratory, and office staff are all very invested in your care. This handbook was designed with your needs in mind to use as a reference and to encourage you to participate in your own care.

IMPORTANT PHONE NUMBERS

• Farmington Office: 844-467-3483 or 860-321-7082

• Hartford Office: 860-525-8283

Note: We have 2 clinical offices where we perform IUIs – both Farmington and Hartford. The Hartford office is closed on weekends and major holidays. All cycling ultrasounds, intrauterine inseminations, IVF procedures, and urgent/emergency care needs are performed ONLY at the Farmington office on weekends and major holidays. (Our New London office has limited ultrasound monitoring hours Monday through Friday and bloodwork. No IUIs are performed in the New London office.)

FOR AFTER HOUR AND WEEKEND EMERGENCIES: BOTH FARMINGTON AND HARTFORD OFFICES:

If you have an emergency and it is after office hours or on weekends, call 844-467-3483 or 860-321-7082 and ask for the physician on call.

If it is a question or concern that can be answered during normal business hours, please leave a message on the nurses' line and we will get back to you when the office re-opens.

"CLINICAL NURSES" LINE:

While you are preparing to cycle (that is, prior to actually starting your injections), call your clinical nurse for any questions or concerns you may have:

For Dr. Nulsen's patients 844-467-3483 X 8026 (Deb J. in

Farmington)

For Dr. Benadiva's patients 844-467-3483 X 8028 (Glori in

Farmington)

For Dr. Schmidt's patients 844-467-3483 X 8035 (Jenny in

Farmington)

For Dr. Engmann's Farmington patients 844-467-3483 X 8035 (Jenny in

Farmington)

For Dr. Engmann's Hartford patients 860-525- 8283 X 15 (Darlene in Hartford) For Drs DiLuigi's patients 860-525- 8283 X 33 (Marie in Hartford)

• Do not leave an emergency message on this line. If your call is not urgent you may not hear from the clinical team until the following business day.

 Messages are not retrieved from these lines after hours or on weekends. However, on Mondays all calls received on the weekend will be retrieved.

ASSISTED REPRODUCTIVE TECHNOLOGIES (ARTS/IVF) NURSES" LINE:

Once you are actually cycling on medications, that is, taking the injections, your care is coordinated by the "ARTs / IVF" Team:

844-467-3483 X 8068 for FARMINGTON OFFICE PATIENTS OR

(860) 525- 8283 X 15 (Dr Engman's Hartford patients) or X 33 (Dr Diluigi's patients) for HARTFORD OFFICE PATIENTS

- Do not leave an emergency message on this line. If your call is not urgent you may not hear from the ARTs/IVF team until the following day.
- Messages are not retrieved from these lines after hours or on weekends. However, on Mondays all calls received on the weekend will be retrieved.

IMPORTANT INSURANCE INFORMATION

Insurance company benefits vary greatly when it comes to infertility treatments. It is your responsibility to discuss benefit coverage with your Financial Service Representative (FSR) here at The Center BEFORE you begin your treatment plan.

If you begin your treatment without completing the pre-certification/financial clearence, you may be required to pay "out of pocket" for procedures, ultrasounds, lab services, blood work, and medications associated with your treatment.

A start date from the medical staff does not mean that you are financially cleared. You must check with your FSR before beginning or making a change to your treatment plan.

Before beginning treatment:

Dr. Nulsen's patients:

Dr. Benadiva's patients:

Dr. Schmidt's patients:

Dr. Schmidt's patients:

Dr. Engmann's FARMINGTON patients:

contact your FSR Susan @ 844-467-3483 X 8015

contact your FSR Veronica @ 844-467-3483 X 8031

contact your FSR Veronica @ 844-467-3483 X 8031

Drs Engmann in HARTFORD contact your FSR Janette @ (860-525-

8283 X 1

Dr. DiLuigi in HARTFORD contact your FSR Janette @ (860-525-

8283 X 1

IMPORTANT PRESCRIPTION MEDICATION INFORMATION

Have your prescriptions filled by 1 week prior to your anticipated cycle start. Infertility medications are specialty items. This is why we have listed pharmacies that specialize in these medications. Your prescription will come with refills so you can call your pharmacy when necessary to activate the refills. Monitor your available medications so that you can order refills with enough lead time for delivery. Be prepared for the weekends when it is more difficult to obtain refills. It is your responsibility to activate the re-fills.

• Some insurance companies are specific and dictate which pharmacy you must go to for coverage. As a result, in certain circumstances, it may take you several weeks to obtain your medications.

Beacon (Southington)	(860) 628-3972	Monday- Saturday
Village Pharmacy	877-334-1610	24 Hours (no Sunday delivery)
Freedom Drug	800-660-4283	24 Hours (no Sunday delivery)
IVP	800-483-8001	24 Hours (no Sunday delivery)
Schrafts	800-876-4545	24 Hours (no Sunday delivery)
Oranoque (Stratford)	888-556-3313	24 Hours (no Sunday delivery)
Apothecary by Design	877-814-8447	24 Hours (no Sunday delivery)
Aetna Specialty	866-782-2779	(insurance specific)
Caremark	877-269-4831 or	800-378-5697 (option 2) (insurance specific)
Tel-Drug (Cigna)	800-351-3606	(insurance specific)
Precision Rx (Anthem)	888-662-0944	(insurance specific)
Medco	1-888-327-9791	(insurance specific)

EMERGENCY MEDICATIONS:

Walgreens Pharmacy
102 Washington St.

New Britain, CT

24 Hour pharmacy
860-826-7272

We can also fax your prescription to any of these pharmacies as per your request.

PRENATAL VITAMINS/FOLIC ACID;

Begin taking prenatal vitamins before you start cycling. Several studies have shown that folic acid supplementation can significantly reduce the occurrence of neural tube defects in infants (such as spina bifida). This can be accomplished by taking the prenatal vitamin we have included in your prescription. This prescription contains 1 mg of folic acid.

NON-PRESCRIPTION MEDICATIONS

Please check with your primary care physician before taking any medications during your cycle.

Some herbal medications (St. John's Wart, Gingko, and Echinacea) have been found to be detrimental to eggs and sperm interaction. UConn Health Center does provide a "Pregnancy Riskline" to answer your questions related to medication and other exposures during pregnancy. You can contact the experts at Mother to Baby @866-626-6847 or online at mothertobaby.org.

CAFFEINE AND SMOKING

Eliminate cigarettes and minimize your intake of caffeine. Both of these substances have been shown to affect fertility. Switch to decaffeinated beverages if needed.

CONSENTS

- Prior to beginning your ovulation induction therapy, your current SIGNED CONSENTS MUST BE IN YOUR MEDICAL RECORD. SEE ADDENDUMS.
- Consents must be witnessed by an employee of the center OR may be notarized. You
 may come in to the office Monday through Friday between 8 AM and 4:30 PM to
 have your consents witnessed OR use a notary at your convenience. SEE
 ADDENDUMS.

ANSWERING MACHINE OR VOICE MAIL

Please have a functioning answering machine or voice mail with either your first name or your phone number identified in the greeting so that we can leave a message on the answering machine if necessary. Check your answering machine/voice mail every day when you are cycling.

www.uconnfertility.com

- This is The Center's own website which contains links to educational programs.
- Click on to "Patient Information" then "Patient Teaching" and you will see a link to Village Pharmacy and Freedom Fertility Pharmacy icons which have video and audio presentations on all medications and injection techniques. Specific instructions for these are your the Medication Checklist (See addendum at end of handbook)
- You will be given a list of your medications to review.

PART 2 – STEPS INVOLVED IN A CYCLE

THE TYPICAL SEQUENCE OF EVENTS INVOLVED IN AN OVULATION INDUCTION CYCLE CONSISTS OF:

1. TAKING INJECTABLE MEDICATIONS

These medications supplement the body's production of the natural hormone FSH. This results in the development and maturation of multiple eggs during the woman's "stimulated" menstrual cycle increasing the probability of pregnancy. Typically a woman takes these medications for approximately 8 to 10 days.

2. A TIMED, MIDCYCLE INJECTION of "HCG" MEDICATION A timed, midcyle, single injection of HCG hormone mimics the "LH surge" which is experienced during a normal menstrual cycle. This results in the final maturation and release of the eggs from the ovaries.

3. A TIMED INTRAUTERINE INSEMINATION (IUI) A timed intrauterine insemination (IUI). This brings the sperm in closer proximity to the eggs and increases the woman's probability of pregnancy

4. ADDING PROGESTERONE

The addition of Progesterone in the later half of the cycle supports the development of the uterine lining to nourish an embryo.

PART 3 GETTING STARTED WITH YOUR CYCLE

1. We monitor your cycle with ultrasounds and blood work to evaluate your response to the medications.

This is done intentionally so that we can help you get in and out as quickly as possible to avoid interrupting your work and personal schedules. Please be assured that you will get the time you need to address your personal needs. Each day your physician and the team will review your blood work and ultrasound results later in the day. Much of this is done by your own physician behind the scenes. If needed, you will receive a call from us to adjust or confirm you treatment plan.

FARMINGTON PATIENT APPOINTMENTS:

• Cycling ultrasounds are done between 7:30 AM and 8:45 AM in Farmington on the second floor. Ultrasounds are on a first come first serve basis. However, please notify us that you will be coming in so that we can prepare your paperwork.

HARTFORD PATIENT APPOINTMENTS:

- Cycling ultrasounds are scheduled in the Hartford office from 7:45 am to approximately 9:00 am. Please call for an appointment time.
- 2. Cycle Day #1 refers to the first day of "full flow" of your menstrual cycle. Any time before midnight that your usual menses (period) starts with your usual "full flow" (not spotting) is called CYCLE DAY # 1
- 3. YOU NEED TO COME IN FOR A BASELINE ULTRASOUND AND BLOOD WORK ON CYCLE DAY #2 OR # 3 BEFORE YOU START MEDICATIONS to be checked to be sure that it is safe to start your medications
- 4. Injectable medications are started on CYCLE DAY #2 or #3 after you have been given the medical OK to start by the clinical team
- 5. To schedule your first ultrasound and blood work appointment follow these instructions:

PATIENT INSTRUCTIONS FOR MAKING ULTRASOUND AND BLOOD WORK APPOINTMENTS

Make your life easier by following these simple instructions:

Monday through Friday:

• If your menses (period) starts during the workday while our office is open (until 4:30 PM), please call your specific physician's office Farmington 844-467-3483 or 860-321-7082 or Hartford @ 860-525-8283 to schedule your baseline appointment for the next day.

Weekends: Special Instructions

- If your menses starts on a Friday after our office is closed (4:30 PM), please come into the Farmington office at 7:30-8:45 AM on Sunday morning for your baseline ultrasound and blood work.
- If your menses starts on a Saturday before 4:30 PM, please come into the Farmington office on Sunday at 7:30-8:45 AM for a baseline ultrasound and blood work.
- If your menses starts after 4:30 PM on a Saturday, please come into your specific physician's office for a baseline ultrasound and blood work on *Monday* morning at 7:30-8:45 AM.
- If your menses starts on a Sunday before 4:30 PM, please come into your specific physician's office on *Monday* morning at 7:30-8:45 AM for a baseline ultrasound and blood work.
- If your menses starts on Sunday after 4:30 PM, please call your specific physician's office on *Monday* and schedule an "appointment" for an ultrasound and blood work on *Tuesday* morning.

GETTING BLOODWORK:

- It is imperative that your blood be drawn on the day it is ordered, at the Center's blood drawing lab, and in a timely manner. We must have your results back early enough in the day (preferable by noon) so that we can evaluate results, make decisions, and communicate those decisions to you that same day. This is a critical component of your cycle.
- During your cycle your lab work <u>MUST BE DRAWN EARLY IN THE MORNING</u>
 <u>BY 9:00AM</u> to ensure that results are available in a timely manner for review at the team rounds.
- The Center's Lab is the Preferred Lab for Patients. All cycling patients MUST have their cycling and follow-up bloods drawn at our offices in Farmington and/or Hartford/New London. All labs have different assays and normal/abnormal ranges and it is important to have consistency. Our team is familiar with these lab assays and ranges and is most confident when making decisions. Furthermore, we receive your results without delay. Your blood must be drawn by 9:00 am to get timely results for your MD to review.
 - A lab requisition is filled out by our clinical staff and will be held for you at our office.
 - O Plan to leave us a phone # at which we may reach you later that afternoon. We may need to call you to adjust your medication plan.

*** ALL WEEKEND CYCLING LABWORK MUST BE DONE AT OUR FARMINGTON OFFICE ONLY BETWEEN 7:30 AM and 9:00AM****

• Farmington Lab Hours

Monday- Friday 7:00 – 9:00 AM Sat & Sunday 7:30 – 9:00 AM

• Hartford lab Hours:

Monday – Friday 7:30 – 9:00 AM

Sat & Sunday 7:30 – 9:00 AM USE Batterson Park Road, Farmington office

 Quest and Community Hospitals are <u>not acceptable</u> for cycling and follow-up blood work unless prior arrangements have been made with our staff for extreme circumstances. Assays may differ between labs interfering with appropriate interpretation and comparison of results for decision making. Community hospitals generally do not have a fast enough turn around time for us to appropriately monitor your cycle.

RESULTS AND CALL BACKS:

Blood work results are not available to us until the afternoon and are then reviewed by the medical team. If we did not give you instructions at your morning ultrasound, then we will call you in the afternoon with your plan. Expect to hear from us between 1:30 PM and 4:00 PM. If you do not hear from us by 4:00 PM, please call the office.

POTENTIAL RISKS AND SIDE EFFECTS ASSOCIATED WITH STIMULATION MEDICATIONS

- Monitoring during your stimulation phase is necessary to help prevent untoward side effects from your medications such as "Ovarian Hyperstimulation".
- Hyperstimulation may result from too exuberant a response by your ovaries to your stimulation medications.
- If monitoring indicates that your ovaries are responding too exuberantly to your
 medications, your medication dosage may be adjusted, or you may be counseled
 about canceling your cycle if you are at increased risk for significant
 HYPERSTIMULATION or High Order MULTIPLE PREGNANCIES (triplets or
 greater).

1. MEDICATION SIDE EFFECTS

o side effects which may be observed during a medication cycle are breast tenderness, mood swings, bloating, fatigue and pelvic discomfort related to ovarian enlargement. Some women form cysts on their ovaries after ovulation. These cysts do not usually cause problems other than pelvic fullness. Sometimes they grow excessively (size of an orange) and may cause abdominal pain or bloating. The resulting condition is referred to as HYPERSTIMULATION.

Significant hyperstimulation, however, is a relatively infrequent occurrence in IUI cycles since relatively small amounts of ovarian stimulation medication are used (compared to an "IVF" In Vitro Fertilization cycle for example). Typically, if hyperstimulation symptoms occur, symptoms generally occur about 1 week after your HCG trigger injection.

Please call us if you develop any problems during this time particularly if you experience dizziness, decreased urination, shortness of breath, severe abdominal pain, or weight gain of more than 3 pounds. These cysts usually start to resolve by the 12th day after your injection of "HCG" unless you conceive. With pregnancy, the symptoms of HYPERSTIMULATION may last a little longer.

Hyperstimulation treatment usually consists of bedrest and increased fluids, especially Gatorade, Powerade, V-8 juice, or canned soups which all contain fluid and electrolytes.

2. MULTIPLE BIRTHS

• Stimulation drugs are associated with an increased risk of multiple birth. The incidence of multiple birth is approximately 20%, usually twins. About 5% of the total pregnancies result in triplets or more. Additionally there may be a slight increase in the rate of miscarriage.

3. OVARIAN CANCER

Several things contribute to a woman's risk of developing ovarian cancer. These include family history of ovarian cancer, personal history of other cancers, and reproductive history. Not experiencing pregnancy in itself is associated with an increased risk of developing ovarian cancer. The more times a woman ovulates in her lifetime, the higher her risk of developing ovarian cancer. For example, a woman who experienced her first menstrual period at a young age, never has a pregnancy, and undergoes menopause at a late age has a higher risk of ovarian cancer than a woman who had a late first period, had several pregnancies and breastfed afterwards, then an early menopause. Birth control pills are protective against ovarian cancer because they prevent ovulation. Most studies have found that ovulation medications do not increase the risk of either ovarian or breast cancer. At this time, there is no definitive evidence to prove that a relationship exists between infertility medications and ovarian cancer. The American Society for Reproductive Medicine came out with this statement in 2002: "More recent research indicates that patients taking ovulation-inducing drugs face no greater cancer risk than the general population". Although future studies could find an association, at this time ovulation-inducing medications appear to be safe for short-term use and do not appear to cause ovarian cancer.

ABOUT SUPPRESSION WITH GANIRELIX ACETATE or CETROTIDE

- On occasion, some women may benefit from a medication which helps to prevent the body from prematurely releasing a naturally occurring hormone "LH". LH allows the egg to burst forth from the follicle. If this happens too early, the eggs may not be mature enough to be successfully fertilized by sperm.
- Medications, marketed under such names as GANIRELIX ACETATE and CETROTIDE, can suppress the body's normal LH surge and prevent the eggs from being released too early.
- If this medication applies to you, we will indicate this to you specifically in your plan and please see the following instructions:
 - Ganirelix Acetate comes in a prefilled syringe that is ready to administer.
 - Cetrotide comes in a ready to mix "kit"
 - You will be given specific instructions on when to start these
 medications. It depends on your cycle day, the size of your follicles
 and/or your estrogen blood work results. Generally, this
 medication is started about half way through your cycle.
 - Once you start cycling, carry one of these prefilled syringes or kits in your purse with you. This way you will be ready to take the medication when directed to do so
 - These medications are typically administered in the MORNING
 - The medication is a "short" needle injection and is given in the abdomen or outer thigh area.
 - If applicable, review the instructions for the administration of this medication on our website

ABOUT YOUR HCG "TRIGGER" MEDICATION

- When both your blood work and ultrasound results indicate that the eggs should be mature, you will be instructed to stop your stimulation medications and instead use an injectable medication that evening called "HCG" (Human Chorionic Gonadotrophin). "HCG" is taken to "trigger" the release of the eggs. Brand names for this same medication include Novarel and Pregnyl.
- "HCG" causes the final maturation of the eggs and the release of the eggs from the ovaries for fertilization.
- "HCG" MUST BE TAKEN AT THE PRESCRIBED TIME to coordinate with your insemination. Generally this time is between 8:00 PM and 10:00 PM
- HCG is generally prescribed in a 10,000 unit dose. The dose may at times be adjusted. If this is the case, we will give you special instructions at the time.
- HCG is a single dose medication and is given with a short needle (subcutaneously) into the abdomen or outer thigh. (Note: the HCG box indicates intramuscular use (long needle injection) but now the accepted practice is to give it as a subcutaneous (short needle) injection.)

- The day you are instructed to take your "HCG", you will also be given an appointment for your partner to produce a sperm sample to be processed for your IUI.
- If you live within a one hour drive from The Center, your partner can produce the sample at home and drop it off at The Center between 7:30 AM and 8:00 AM. The sample should be protected from heat and cold. A suggestion is to transport the container tucked inside his shirt so that body heat maintains the sample. We will give you the container for this. <u>Your partner will need to drop off the sample and bring a photo ID (driver's license) with him.</u>
- Note: If your partner cannot drop off the sample himself in the morning, then you may strongly wish to have him arrange an appointment here at the center to produce a sample for cryopreservation (freezing). This way the sample will be available to you if such a situation arises. Call 844-467-3483 or 860-321-7082(Farmington) or 860-525-8283 (Hartford) to make an appointment for this.
- If your partner will produce the sample here in the office, he will need to be here at 8:00 AM. There are 3 "production" rooms at The Center in Farmington and one in Hartford. Therefore, there may be a delay in room availability depending upon the particular day.
- If your partner's sperm is being used for the insemination (as opposed to donor sperm), it would be ideal if he abstains from ejaculating for at least 2 days, but not more than 5 days prior to the insemination.
- Processing the sample takes approximately 2 hours. You will be asked to return to The Center for your IUI procedure at 11:00 AM.

ABOUT YOUR INTRAUTERINE INSEMINATION (IUI)

- An intrauterine insemination is a procedure which allows sperm to be deposited directly inside the uterus.
- The purpose of the IUI is to increase the woman's ability to become pregnant by both washing the sperm to enhance its motility and by mechanically placing the sperm closer to the egg.
- An intrauterine insemination is done by using a small flexible catheter.
- You will be positioned as if you were having a PAP test. A speculum is inserted into the vagina and using a very thin flexible catheter, the washed sperm is drawn up into a syringe and gently guided through the cervical opening of the uterus and into the body of the uterus. Minimal cramping, as experienced during a pap test, may be experienced by some women.
- The procedure typically takes a few minutes. You will be asked to rest for about 5 to 10 minutes after the procedure before going back to home or work.
- You can resume your normal activities (including intercourse) unless counseled otherwise by your physician.

("CRINONE" GEL or "ENDOMETRIN" or PROGESTERONE SUPPOSITORIES)

- Progesterone supports the uterine lining and prepares a nourishing environment for the embryo
- Progesterone comes in various formulations including: vaginal gel applicators called "Crinone"; vaginal ovules called "Endometrin"; and natural progesterone vaginal suppositories
- Progesterone is added in to your cycle beginning the SECOND MORNING AFTER vour IUI
- You will stay on progesterone until the onset of your next period (in the event that you are not pregnant) or until instructed to discontinue the medication at your pregnancy ultrasound appointment (approximately 4 to 5 weeks after your IUI)
- See "Helpful Hints for Progesterone" ADDENDUM at the end of this handbook for administration tips for these progesterone supplements.

POST INSEMINATION FOLLOW-UP INSTRUCTIONS

- If you physician has specific instructions regarding follow-up for your cycle, these instructions will be reviewed with you.
- If you do not have a period by 16 days after your IUI, please call us at our office (excluding weekends) and leave a message on the clinical line Farmington 844-467-3483 or 860-321-7082; Hartford office 860-525-8283 X 15 or X 33. (Sometimes your period can be slightly delayed when using progesterone.) Include your name, spelling of your name, (or date of birth) and # at which we can reach you. We will call you back with instructions during scheduled business hours.
- If your pregnancy test is positive, you will receive instructions for follow-up blood work and we will schedule you for a pregnancy ultrasound for approximately 4 to 5 weeks after your IUI. This is a transvaginal ultrasound.

PART 4 TIME TO REVIEW THE TEACHING WEBSITE

- You are responsible for reviewing all assigned medication videos. See MEDICATION CHECKLIST ADDENDUM AT END of handbook
- To reach the website:
 - Log on to www.uconnfertility.com
 - Select "Patient Information"
 - Select "Patient Teaching and Resources"
 - CLICK on "Village Pharmacy" icon
 - Click on to each assigned medication(s)

OR

• CLICK on Freedom Fertility Pharmacy icon and select your medication and injection videos from the list

• Once you have reviewed both this teaching handbook AND the medication and injection videos, call your clinical nurse and leave a message at:

Farmington Office 844-467-3483 or 860-321-7082 (see page 4 for your nurse's phone extension)

Hartford Office 860-525 – 8283 (see page 4 for your nurse's phone extension)

• Your call will be returned WITHIN A FEW DAYS to complete a short phone interview to verify your understanding of your current medication protocol; to answer any questions that you might have; and to receive your actual prescription.

SUMMARY & KEY POINTS TO REMEMBER

- 1. You will not be able to start a cycle until you have financial and medical clearances
- 2. Signed consents MUST be on your medical record before you begin to cycle.
- 3. Call the first day of your period to schedule your baseline ultrasound and blood work as possible.
- 4. Typically injectable medications will start on Cycle day #2 or #3
- 5. You will be told when to start your injectable medications at your baseline ultrasound
- 6. Typically, you will be on these injectable medications for approximately 8 to 10 days
- 7. During this time your progress will be monitored with blood work and ultrasounds until your follicles/eggs are matured
- 8. You will be told when to take the HCG "trigger" injection
- 9. You will be scheduled for an Intrauterine Insemination (IUI) typically 2 days after the HCG "trigger"injection)
- 10. You will begin progesterone for uterine support the *second* morning after your insemination

Please remember that we are here to assist you in any way that we can! Farmington Office # 844-467-3483 or 860-321-7082 Hartford Office # (860) 525-8283

ADDENDUMS

HELPFUL HINTS WHEN TAKING INJECTABLE MEDICATIONS:

- Take the medications at approximately the same time of day (within 1 hour leeway). Generally, evening medications are given between 6:00 PM to 8:00 PM. If your medication plan calls for injections twice a day, space them about 12 hours apart.
- Before injecting the medication, apply ice (wrapped in a cloth) for a few minutes to the area where the medication will be given to help minimize discomfort
- "Rotate" your injection sites for greater comfort
- To avoid bruising at the injection site, apply pressure with a dry gauze pad for approximately one minute to the site

ABOUT MENOPUR INJECTIONS:

• Based upon our patients' experience, you may find it easier to mix WITHOUT THE Q CAP. It is really a matter of preference. ATTACHED PLEASE SEE A VISUAL INSTRUCTION ADDENDUM FOR MIXING MENOPUR WITH A NEEDLE and SYRINGE (WITHOUT the "Q CAP" adaptor that comes in the box).

HELPFUL HINTS FOR USING CRINONE (PROGESTERONE) GEL:

• Insert Crinone ONCE A DAY in the morning into the vagina so that the gel will spread evenly throughout the day with normal daily activities. This will help minimize "clumping" of the gel in the vagina. Insert the applicator approximately one third of the depth of the applicator length into the vagina.

HELPFUL HINTS FOR ENDOMTRIN (PROGESTERONE) OVULES:

• The usual dose for endomterin ovules is 1 ovule TWICE A DAY (morning and evening) inserted into the vagina with the endometrin applicator.

HELPFUL HINTS FOR USING PROGESTERONE SUPPOSITORIES:

- Vaginal suppositories are inserted into the vagina TWICE A DAY once in the morning and then again in the evening
- Insert your morning suppository approximately ½ hour before you get up out of bed
- Insert suppository approximately 2/3 the length of your index finger (Inserting the progesterone suppository too far into the vagina and up to the cervix may cause some "spotting" to occur.)
- Wear a mini-pad for greater comfort and to protect your clothing
- Intercourse is permitted when on progesterone suppository therapy. However, refrain from having intercourse for approximately ½ hour after inserting the suppository to allow it to melt.

Patient ID#

Medication Checklist (Revised 8.15)

_Cetrotide0.25 mg3mg AND "Subcutaneous injection" videos
Ganirelix Acetate 250 mcg AND "Subcutaneous injection" videos
Leuprolide Acetate(2 week kit) AND "Subcutaneous injection" videos
Microdose Lupron (40mcg/0.1ml) AND "Subcutaneous injection" videos
Follistim pen Dose:
_Gonal-F pen Dose:
 Luveris vial (75IU) AND "Subcutaneous injection" videos
 Menopur (75IU) (without Q cap) AND "Subcutaneous injection videos
*See the addendum attached in packet for hints for mixing without the Q cap
 _Repronex (75IU) (without Q cap) and Subcutaneous injection" videos
 _Ovidrel 250 mcg AND "Subcutaneous injection" videos
 _HCG (10,000IU) AND "Subcutaneous injection" videos
 _Leuprolide Acetate 0.5cc (1mg/0.2ccvial) AND "Subcutaneous injection"
videos
 Progesterone in sesame oil (50mg/cc) AND "Intramuscular injection" videos
 Progesterone in olive oil (50mg/cc) AND "Intramuscular injection" videos
 Progesterone in ethyl oleate (50mg/cc) AND "Intramuscular injection"
videos
 _Crinone 8% gel vaginal applicator
 _Endometrin vag insert (100mg)
 _Progesterone supp 50mg 100mg
 _Vivelle dots (0.1mg)
 _Bravelle (75IU) AND "Subcutaneous injection" videos

Please review all medications and injection techniques checked above on the online video injection lesson at www.uconnfertility.com

- Next choose "Patient Information"
- Then "Patient Teaching and resources"
- Click on Village Fertility Pharmacy icon
- Click on to each assigned medications and watch the medication video and injection technique
- *OR* can click on <u>Freedom Fertility Pharmacy</u> icon and bring up list of medications and injections to review.
- Both sites are in English and Spanish